

Modeling misspecification as a parameter in Bayesian structural equation models

James Ohisei Uanhoro
james.uanhoro@unt.edu

May 4, 2023

Department of Educational Psychology
University of North Texas

Outline

Introduction

Model

Simulation studies

Closing

Introduction

Misspecification is active research area for Bayesian SEMs

- Levy (2011): Posterior predictive checking using the likelihood ratio test; Bayesian SRMR
- Hoofs, van de Schoot, Jansen, and Kant (2018): Bayesian RMSEA
- Garnier-Villarreal and Jorgensen (2020): Several translations of frequentist fit indices: RMSEA, CFI, TLI, ...
- Cain and Zhang (2019): Deviance information criterion
- Fit hypothesized model $\xrightarrow{\text{then}}$ compute misspecification

Fit hypothesized model and estimate misspecification simultaneously

- Wu and Browne (2015) in frequentist context:
 - Parameter estimates have greater uncertainty about them reflecting degree of model misspecification
 - Estimates the RMSEA as a model parameter
- We could simply recreate Wu and Browne (2015) in a Bayesian context, but:
 - The RMSEA is not so easy to interpret (e.g. Chen, Curran, Bollen, Kirby, & Paxton, 2008; Savalei, 2012)
 - The SRMR has clearer interpretations.
 - Model misspecification so parameter estimates reflect degree of model incorrectness, while estimating an SRMR-type index.

Fit hypothesized model and estimate misspecification simultaneously

- Wu and Browne (2015) in frequentist context:
 - Parameter estimates have greater uncertainty about them reflecting degree of model misspecification
 - Estimates the RMSEA as a model parameter
- We could simply recreate Wu and Browne (2015) in a Bayesian context, but:
 - The RMSEA is not so easy to interpret (e.g. Chen et al., 2008; Savalei, 2012)
 - The SRMR has clearer interpretations.
 - Model misspecification so parameter estimates reflect degree of model incorrectness, while estimating an SRMR-type index.

Fit hypothesized model and estimate misspecification simultaneously

- Wu and Browne (2015) in frequentist context:
 - Parameter estimates have greater uncertainty about them reflecting degree of model misspecification
 - Estimates the RMSEA as a model parameter
- We could simply recreate Wu and Browne (2015) in a Bayesian context, but:
 - The RMSEA is not so easy to interpret (e.g. Chen et al., 2008; Savalei, 2012)
 - The SRMR has clearer interpretations.
 - Model misspecification so parameter estimates reflect degree of model incorrectness, while estimating an SRMR-type index.

Fit hypothesized model and estimate misspecification simultaneously

- Wu and Browne (2015) in frequentist context:
 - Parameter estimates have greater uncertainty about them reflecting degree of model misspecification
 - Estimates the RMSEA as a model parameter
- We could simply recreate Wu and Browne (2015) in a Bayesian context, but:
 - The RMSEA is not so easy to interpret (e.g. Chen et al., 2008; Savalei, 2012)
 - The SRMR has clearer interpretations.
 - Model misspecification so parameter estimates reflect degree of model incorrectness, while estimating an SRMR-type index.

Fit hypothesized model and estimate misspecification simultaneously

- Wu and Browne (2015) in frequentist context:
 - Parameter estimates have greater uncertainty about them reflecting degree of model misspecification
 - Estimates the RMSEA as a model parameter
- We could simply recreate Wu and Browne (2015) in a Bayesian context, but:
 - The RMSEA is not so easy to interpret (e.g. Chen et al., 2008; Savalei, 2012)
 - The SRMR has clearer interpretations.
 - Model misspecification so parameter estimates reflect degree of model incorrectness, while estimating an SRMR-type index.

Fit hypothesized model and estimate misspecification simultaneously

- Wu and Browne (2015) in frequentist context:
 - Parameter estimates have greater uncertainty about them reflecting degree of model misspecification
 - Estimates the RMSEA as a model parameter
- We could simply recreate Wu and Browne (2015) in a Bayesian context, but:
 - The RMSEA is not so easy to interpret (e.g. Chen et al., 2008; Savalei, 2012)
 - The SRMR has clearer interpretations.
 - Model misspecification so parameter estimates reflect degree of model incorrectness, while estimating an SRMR-type index.

Fit hypothesized model and estimate misspecification simultaneously

- Wu and Browne (2015) in frequentist context:
 - Parameter estimates have greater uncertainty about them reflecting degree of model misspecification
 - Estimates the RMSEA as a model parameter
- We could simply recreate Wu and Browne (2015) in a Bayesian context, but:
 - The RMSEA is not so easy to interpret (e.g. Chen et al., 2008; Savalei, 2012)
 - The SRMR has clearer interpretations.
 - **Model misspecification so parameter estimates reflect degree of model incorrectness, while estimating an SRMR-type index.**

Model

CFA as an example

Begin with Muthén and Asparouhov (2012):

$$\Sigma = \underbrace{\Lambda\Phi\Lambda^T + \Delta}_{\text{Standard}} + \Psi \quad (1)$$

Λ : Loading matrix, Φ : Interfactor correlation matrix, Δ : Standard residual covariance matrix (often diagonal)

Ψ : Residual covariance matrix, with all off-diagonal elements estimated.

- Theoretically: Assumed to reflect the influence of minor factors (MacCallum & Tucker, 1991)
- Practically: Ψ is not identified. Muthén and Asparouhov (2012) used an inverse-Wishart prior with known parameters to shrink elements in Ψ to zero.

CFA as an example

Begin with Muthén and Asparouhov (2012):

$$\Sigma = \underbrace{\Lambda\Phi\Lambda^T + \Delta}_{\text{Standard}} + \Psi \quad (1)$$

Λ : Loading matrix, Φ : Interfactor correlation matrix, Δ : Standard residual covariance matrix (often diagonal)

Ψ : Residual covariance matrix, with all off-diagonal elements estimated.

- Theoretically: Assumed to reflect the **influence of minor factors** (MacCallum & Tucker, 1991)
- Practically: Ψ is not identified. Muthén and Asparouhov (2012) used an inverse-Wishart prior with known parameters to shrink elements in Ψ to zero.

CFA as an example

Begin with Muthén and Asparouhov (2012):

$$\Sigma = \underbrace{\Lambda\Phi\Lambda^T + \Delta}_{\text{Standard}} + \Psi \quad (1)$$

Λ : Loading matrix, Φ : Interfactor correlation matrix, Δ : Standard residual covariance matrix (often diagonal)

Ψ : Residual covariance matrix, with all off-diagonal elements estimated.

- Theoretically: Assumed to reflect the **influence of minor factors** (MacCallum & Tucker, 1991)
- Practically: Ψ is not identified. Muthén and Asparouhov (2012) used an inverse-Wishart prior with known parameters to shrink elements in Ψ to zero.

Goal is to model misspecification

$$\Sigma = \underbrace{\Lambda \Phi \Lambda^T + \Delta}_{\text{Standard}} + \Psi$$

Let ψ_{ij} ($i \neq j$) be off diagonal elements in Ψ , reflecting misspecification / minor factor influences.

To model minor factor influences:

$$\frac{\psi_{ij}}{\underbrace{\sqrt{\sigma_{jj}\sigma_{ii}}}_{\text{SRCs}}} \sim \mathcal{N}(0, \tau), \quad \tau \sim \mathcal{N}^+(0, 1)$$

σ_{ii}/jj : indicator variances i.e. SRCs: standardized residual covariances.

Goal is to model misspecification

$$\Sigma = \underbrace{\Lambda \Phi \Lambda^T + \Delta}_{\text{Standard}} + \Psi$$

Let ψ_{ij} ($i \neq j$) be off diagonal elements in Ψ , reflecting misspecification / minor factor influences.

To model minor factor influences:

$$\frac{\psi_{ij}}{\underbrace{\sqrt{\sigma_{jj}\sigma_{ii}}}_{\text{SRCs}}} \sim \mathcal{N}(0, \tau), \quad \tau \sim \mathcal{N}^+(0, 1)$$

σ_{ii}/σ_{jj} : indicator variances i.e. SRCs: standardized residual covariances.

A hierarchical model for SRCs

$$\frac{\psi_{ij}}{\underbrace{\sqrt{\sigma_{jj}\sigma_{ii}}}_{\text{SRCs}}} \sim \mathcal{N}(0, \tau), \quad \tau \sim \mathcal{N}^+(0, 1)$$

Implications

- SRCs are assumed to be zero on average
- SRC scale parameter (τ) is the root mean square error of SRCs with location constrained to be zero
- τ is akin to the correlation root mean square residual (CRMR)
- Smaller values of τ reflect a model with smaller SRCs, lower misspecification OR smaller influence of minor factors
- $\tau < 0.05 \implies$ most SRCs < 0.10

A hierarchical model for SRCs

$$\frac{\psi_{ij}}{\underbrace{\sqrt{\sigma_{jj}\sigma_{ii}}}_{\text{SRCs}}} \sim \mathcal{N}(0, \tau), \quad \tau \sim \mathcal{N}^+(0, 1)$$

Implications

- SRCs are assumed to be zero on average
- SRC scale parameter (τ) is the root mean square error of SRCs with location constrained to be zero
- τ is akin to the correlation root mean square residual (CRMR)
- Smaller values of τ reflect a model with smaller SRCs, lower misspecification OR smaller influence of minor factors
- $\tau < 0.05 \implies$ most SRCs < 0.10

A hierarchical model for SRCs

$$\underbrace{\frac{\psi_{ij}}{\sqrt{\sigma_{jj}\sigma_{ii}}}}_{\text{SRCs}} \sim \mathcal{N}(0, \tau), \quad \tau \sim \mathcal{N}^+(0, 1)$$

Implications

- SRCs are assumed to be zero on average
- SRC scale parameter (τ) is the root mean square error of SRCs with location constrained to be zero
- τ is akin to the correlation root mean square residual (CRMR)
- Smaller values of τ reflect a model with smaller SRCs, lower misspecification OR smaller influence of minor factors
- $\tau < 0.05 \implies$ most SRCs < 0.10

Simulation studies

Three simulation studies

List of simulation studies

1. Can the proposed approach recover model parameters?
 - Yes, upcoming
2. Can differences in τ between competing models be used for model selection?
 - Lacks power, more promising to use **approximate leave-one-out cross validation** to select between competing models.
3. Can the Ψ matrix be used to detect specific residual covariances that are too large?
 - Yes, especially for larger samples.

Three simulation studies

List of simulation studies

1. Can the proposed approach recover model parameters?
 - Yes, upcoming
2. Can differences in τ between competing models be used for model selection?
 - Lacks power, more promising to use **approximate leave-one-out cross validation** to select between competing models.
3. Can the Ψ matrix be used to detect specific residual covariances that are too large?
 - Yes, especially for larger samples.

Three simulation studies

List of simulation studies

1. Can the proposed approach recover model parameters?
 - Yes, upcoming
2. Can differences in τ between competing models be used for model selection?
 - Lacks power, more promising to use **approximate leave-one-out cross validation** to select between competing models.
3. Can the Ψ matrix be used to detect specific residual covariances that are too large?
 - Yes, especially for larger samples.

Three simulation studies

List of simulation studies

1. Can the proposed approach recover model parameters?
 - Yes, upcoming
2. Can differences in τ between competing models be used for model selection?
 - Lacks power, more promising to use **approximate leave-one-out cross validation** to select between competing models.
3. Can the Ψ matrix be used to detect specific residual covariances that are too large?
 - Yes, especially for larger samples.

Study 1: Recovery of structural parameters

Simulate data with minor factor influences (Ψ)

5 modeling approaches

- Two **ignore** minor factor influences:
 1. *Freq*: Standard frequentist
 2. *Baseline*: Standard Bayesian
- One **fixes** the size of minor factor influences:
 3. *AZ*: Muthén and Asparouhov (2012)
- Two **estimate** the size of minor factor influences:
 4. *LKJ*: another proposed method (in paper)
 5. *NRM*: hierarchical normal method (focus here)

Results: Bias adequate; Coverage poor for methods that ignore or fix the size of the influence of minor factors

Study 1: Recovery of structural parameters

Simulate data with minor factor influences (Ψ)

5 modeling approaches

- Two **ignore** minor factor influences:
 1. *Freq*: Standard frequentist
 2. *Baseline*: Standard Bayesian
- One **fixes** the size of minor factor influences:
 3. *AZ*: Muthén and Asparouhov (2012)
- Two **estimate** the size of minor factor influences:
 4. *LKJ*: another proposed method (in paper)
 5. *NRM*: hierarchical normal method (focus here)

Results: Bias adequate; Coverage poor for methods that ignore or fix the size of the influence of minor factors

Study 1: Recovery of structural parameters

Simulate data with minor factor influences (Ψ)

5 modeling approaches

- Two **ignore** minor factor influences:
 1. *Freq*: Standard frequentist
 2. *Baseline*: Standard Bayesian
- One **fixes** the size of minor factor influences:
 3. *AZ*: Muthén and Asparouhov (2012)
- Two **estimate** the size of minor factor influences:
 4. *LKJ*: another proposed method (in paper)
 5. *NRM*: hierarchical normal method (focus here)

Results: Bias adequate; Coverage poor for methods that ignore or fix the size of the influence of minor factors

Study 1: Recovery of structural parameters

Simulate data with minor factor influences (Ψ)

5 modeling approaches

- Two **ignore** minor factor influences:
 1. *Freq*: Standard frequentist
 2. *Baseline*: Standard Bayesian
- One **fixes** the size of minor factor influences:
 3. *AZ*: Muthén and Asparouhov (2012)
- Two **estimate** the size of minor factor influences:
 4. *LKJ*: another proposed method (in paper)
 5. *NRM*: hierarchical normal method (focus here)

Results: Bias adequate; Coverage poor for methods that ignore or fix the size of the influence of minor factors

Study 1: Recovery of structural parameters

Simulate data with minor factor influences (Ψ)

5 modeling approaches

- Two **ignore** minor factor influences:
 1. *Freq*: Standard frequentist
 2. *Baseline*: Standard Bayesian
- One **fixes** the size of minor factor influences:
 3. *AZ*: Muthén and Asparouhov (2012)
- Two **estimate** the size of minor factor influences:
 4. *LKJ*: another proposed method (in paper)
 5. *NRM*: hierarchical normal method (focus here)

Results: Bias adequate; Coverage poor for methods that ignore or fix the size of the influence of minor factors

Study 1: Recovery of structural parameters

Simulate data with minor factor influences (Ψ)


5 modeling approaches

- Two **ignore** minor factor influences:
 1. *Freq*: Standard frequentist
 2. *Baseline*: Standard Bayesian
- One **fixes** the size of minor factor influences:
 3. *AZ*: Muthén and Asparouhov (2012)
- Two **estimate** the size of minor factor influences:
 4. *LKJ*: another proposed method (in paper)
 5. *NRM*: hierarchical normal method (focus here)

Results: Bias adequate; Coverage poor for methods that ignore or fix the size of the influence of minor factors

Study 1: Empirical coverage rate of 90% CI

Coverage of structural parameters and τ'




Findings:

- *Freq* and *Baseline* models fail when sample size or τ is large
- Performance of *AZ* model deteriorates at large τ
- Methods that estimate the size of minor factor influences perform as expected

An empirical example: Hospital Anxiety and Depression scale

Bifactor model parameters, $n = 21820$, $\tau = 0.028$. Note narrow intervals for the *Standard* approach.



LX.Y are loadings of factor Y on item X; RV: residual variances. Standard estimates are from a model without Ψ .

Closing

Discussion points / future work

- Approach works for fitting hypothesized models simultaneously with misspecification
- There is an R package for this: `minorbsem`
- Package also includes:
 - Implementation of Wu and Browne (2015) method, other priors for standardized residual covariances e.g. lasso, generalized double Pareto (global-local)
 - Global-local prior on all cross-loadings, relaxing simple structure
- Extensions to other data (e.g. binary)
 - Additional issues/extensions are on GitHub:
<https://github.com/jamesuanhoro/minorbsem/issues/3>

References

- Cain, M. K., & Zhang, Z. (2019). Fit for a Bayesian: An evaluation of PPP and DIC for structural equation modeling. *Structural Equation Modeling*, 26(1), 39–50. doi: 10.1080/10705511.2018.1490648
- Chen, F., Curran, P. J., Bollen, K. A., Kirby, J., & Paxton, P. (2008, May). An empirical evaluation of the use of fixed cutoff points in RMSEA test statistic in structural equation models. *Sociological Methods & Research*, 36(4), 462–494. doi: 10.1177/0049124108314720
- Garnier-Villarreal, M., & Jorgensen, T. D. (2020, February). Adapting fit indices for Bayesian structural equation modeling: Comparison to maximum likelihood. *Psychological Methods*, 25(1), 46–70. doi: 10.1037/met0000224
- Hoofs, H., van de Schoot, R., Jansen, N. W. H., & Kant, I. (2018, August). Evaluating model fit in Bayesian confirmatory

factor analysis with large samples: Simulation study introducing the BRMSEA. *Educational and Psychological Measurement*, 78(4), 537–568. doi: 10.1177/0013164417709314

Levy, R. (2011, October). Bayesian data-model fit assessment for structural equation modeling. *Structural Equation Modeling: A Multidisciplinary Journal*, 18(4), 663–685. doi: 10.1080/10705511.2011.607723

MacCallum, R. C., & Tucker, L. R. (1991). Representing sources of error in the common-factor model: Implications for theory and practice. *Psychological Bulletin*, 109, 502–511. doi: 10.1037/0033-2909.109.3.502

Muthén, B. O., & Asparouhov, T. (2012). Bayesian structural equation modeling: A more flexible representation of substantive theory. *Psychological Methods*, 17(3), 313–335.

doi: 10.1037/a0026802

- Savalei, V. (2012, December). The relationship between root mean square error of approximation and model misspecification in confirmatory factor analysis models. *Educational and Psychological Measurement, 72*(6), 910–932. doi: 10.1177/0013164412452564
- Wu, H., & Browne, M. W. (2015, September). Quantifying adventitious error in a covariance structure as a random effect. *Psychometrika, 80*(3), 571–600. doi: 10.1007/s11336-015-9451-3